




# Double burden of underweight and overweight among women of reproductive age in Bangladesh

Raaj Kishore Biswas<sup>1,\*</sup> , Nusma Rahman<sup>2</sup>, Rasheda Khanam<sup>3</sup>, Abdullah H Baqui<sup>3</sup> and Saifuddin Ahmed<sup>4</sup>

<sup>1</sup>Transport and Road Safety (TARS) Research Centre, University of New South Wales, Old Main Building (K15) – Floor 1, Sydney, NSW 2052, Australia: <sup>2</sup>Department of Statistics, Jagannath University, Dhaka, Bangladesh: <sup>3</sup>Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA: <sup>4</sup>Department of Population, Family and Reproductive Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

Submitted 28 March 2019: Final revision received 28 May 2019: Accepted 11 June 2019

## Abstract

**Objective:** Bangladesh, like many emerging economies of South-East Asia, has started to experience a double burden of continuing high rates of undernutrition and increasing rates of overweight and obesity. A lack of assessment of the nutritional shift leaves a gap in current policies: the growing overweight and obesity is yet to be addressed. The present paper investigates the change in nutritional status, particularly the shift in BMI, of Bangladeshi women of reproductive age (15–49 years) and characterizes the vulnerable households for both underweight and overweight status during a period of 10 years (2004–2014).

**Design:** Generalized linear mixed-effect models were fitted for both urban and rural residents to assess underweight and overweight status.

**Setting:** Bangladesh Demographic and Health Surveys.

**Participants:** Women aged 15–49 years ( $n$  53 077).

**Results:** The proportion of overweight increased during 2004–2014 from 10.7 to 25.1% and the proportion of underweight decreased from 32.6 to 18.2%. Prevalence of underweight status remained high in rural areas and prevalence of overweight increased rapidly in both rural and urban areas, creating a double burden. The significant contributors to this double burden were the change in women's level of education, increased household wealth, divisional location and rapid urbanization.

**Conclusions:** The findings indicate that specific cohort- or area-based intervention policy studies in line with the UN Decade of Action on Nutrition are required to address the nutritional double burden in Bangladesh.

**Keywords**  
Nutrition transition  
BMI  
Overweight  
Obesity  
Underweight  
Bangladesh

In the last 30 years, average BMI has increased annually by about 0.4 kg/m<sup>2</sup> worldwide<sup>(1)</sup>; the increase has occurred primarily in developing countries along with an alarming rate of increase in overweight and obesity rates<sup>(2–4)</sup>. Asian countries are now part of this global trend, where the continuing burden of undernutrition and the emerging problem of overnutrition have created a double burden of malnutrition<sup>(5–7)</sup>. Bangladesh, like many emerging economies of South-East Asia, has started to experience a double burden of continuing high rates of undernutrition and increasing rates of overweight and obesity<sup>(8,9)</sup>.

Bangladesh has been fighting both poverty and undernourishment for a long time and has progressed towards being a low middle-income nation with decreased rates

of undernutrition<sup>(10–14)</sup>. The country has recently shown remarkable improvement in the public health sector, including the achievement of both Millennium Development Goal 4 and Millennium Development Goal 5 ahead of time<sup>(15,16)</sup> and striving to achieve the Sustainable Development Goals<sup>(17–19)</sup>. This was facilitated by a consistent, more than 6.0% annual rate of growth in gross domestic product along with the reduction of poverty by one-third in the last decade<sup>(20,21)</sup>. The policies regarding nutritional status of women, particularly of mothers, are focused on undernutrition and underweight<sup>(22–24)</sup>. However, a lack of assessment of the nutritional shift leaves a gap in current policies: the growing overweight and obesity is yet to be addressed<sup>(25–27)</sup>.

\*Corresponding author: Email raajkishore.biswas@student.unsw.edu.au

Low BMI has a significant association with preterm birth and other adverse birth outcomes. Women in developing countries generally face problems of underweight and undernourishment, to which poverty, food scarcity and illiteracy directly contribute<sup>(28–31)</sup>. Underweight (BMI < 18.50 kg/m<sup>2</sup>) during the reproductive age of women is associated with increased risk of low birth weight, preterm birth and intra-uterine growth restriction, as well as neonatal morbidity, mortality and growth failure<sup>(32–35)</sup>. However, an improvement in the nutritional status and a decreasing trend of underweight are evident in sub-Saharan Africa and South Asia<sup>(36,37)</sup>. Microfinance programmes, growth in gross domestic product, better diet, accessibility to health care and increased female education have contributed to the positive nutritional shift in Bangladesh, which decreased rates of underweight status<sup>(31,38)</sup>.

Bangladesh is experiencing a new health problem of overweight and obesity, which are major risk factors for non-communicable diseases, especially diabetes and CVD, and are estimated to cause 3.8% of the disability-adjusted life-years worldwide<sup>(39–42)</sup>. Rapid urbanization, decreased physical activity, change in lifestyle and change in dietary intake are the major contributors to this shift in low- and middle-income countries such as Bangladesh<sup>(43–46)</sup>. A policy-based discussion was initiated by Khan and Talukder in 2013<sup>(8)</sup> regarding the double burden of malnutrition in Bangladesh, where they identified the insufficient investment to address overweight and obesity in Bangladesh along with the lack of appropriate health policies.

The present paper investigates the change in nutritional status, particularly the shift in BMI, of Bangladeshi women of reproductive age (15–49 years). With the current policies focusing on undernutrition in rural areas, Bangladesh may have overlooked overweight and obesity, which is expected to be more prominent in urban affluent areas. Therefore the paper further assesses the gap between rural and urban nutritional status to characterize the vulnerable households for both underweight and overweight status for women in Bangladesh. The trend of BMI over a period of 10 years (2004–2014) was analysed using the last four rounds of Bangladesh Demographic and Health Surveys (BDHS) data. These should contribute to reassess the existing nutritional health policies in the region.

## Materials and methods

### Data description

The Demographic and Health Surveys (DHS) collect data from more than ninety low- and middle-income countries on the demographic and health characteristics of populations approximately at 5-year intervals<sup>(47)</sup>. The BDHS have been conducted among ever-married women of reproductive age (15–49 years) since 1993<sup>(15)</sup>. These nationally representative cross-sectional surveys are conducted in

collaboration with the National Institute of Population Research and Training, Bangladesh, ICF International, USA, and Mitra and Associates, Bangladesh.

Two-stage stratified cluster sampling techniques were adapted for sampling purposes in these surveys<sup>(15)</sup>. The sampling frame for the survey was a complete list of enumeration areas from the most recent census, which were either a village (*mouza* in rural area and *moballa* in urban area) or a part of a village or a group of villages. In the first stage of the sampling, the enumeration areas (clusters) were selected using the probability-proportional-to-size sampling method, where 600 clusters were selected in all the BDHS. In the second stage, an equal-probability systematic sampling method was applied to draw an average of thirty households from each selected cluster. The data for the current study were extracted from the BDHS conducted in 2004, 2007, 2011 and 2014. Only the data of the female respondents were considered; temporary residents (*de jure v. de facto*) and pregnant women were excluded from the analytic data set. Over 10 000 women were considered as the final sample size for each survey and they were merged together for the current study, constituting an overall sample of size 53 077.

### Outcome variable

The present study applied the typical BMI categories constructed by the WHO: <18.50 kg/m<sup>2</sup> = underweight, 18.50–24.99 kg/m<sup>2</sup> = normal, ≥25.00–29.99 kg/m<sup>2</sup> = overweight and ≥30.00 kg/m<sup>2</sup> = obese<sup>(48)</sup>. Continuous scaling of BMI at conservative form was used here to ensure all weight/height<sup>2</sup> were captured. As the number of obese women was about 0.1%, those obese were combined with the overweight category of BMI. While analysing the underweight level, a dichotomous variable of underweight *v.* others (normal/overweight/obese) was created. A similar process was followed for assessing overweight (BMI ≥ 25.00 kg/m<sup>2</sup>) *v.* others (normal/underweight).

The present paper discusses the disproportional burden of nutrition in two residencies: urban and rural. Although the variable 'residency' was taken from the BDHS, it should be noted that DHS surveys never define the urban–rural in a surveyed country, rather they use the country-specific administrative definition. According to the Bangladesh Bureau of Statistics, urban areas are comprised of 'a central place having 5000 population with such amenities as metalled roads, improved communication, electricity, gas, water supply, sewerage, sanitation and also having comparatively higher density of population with majority population in non-agriculture occupations<sup>(49)</sup>'. These includes cities, towns, *Paurasavas* (municipalities) and Cantonments.

### Statistical analysis

BDHS provides a range of household-level socio-demographic factors. Four surveys from 2004 to 2014, each in an interval of 3 years, were used in the present study.



An overview of the variables was conducted through frequency distribution (Table 1), where changes in socio-demographic factors over the years are displayed as well as the categories of BMI. Bivariate analysis between the BMI categories and sociodemographic factors was conducted. Their primary association was evaluated using the  $\chi^2$  test, which was adjusted for the complex survey design (survey weights and cluster-wise variation) using second-order Rao–Scott corrections<sup>(50)</sup>. To assess the trend of the nutritional burden, generalized linear mixed-effect models<sup>(51)</sup> were applied, for both the underweight and the overweight variable. The time variable ‘year’ was considered as a covariate for assessing the trend.

The generalized linear mixed-effect models were fitted separately for urban and rural areas with year as their covariate (Tables 3 and 4). Each model was adjusted for age, parity, education, wealth index, division and marital status. The final model included urban and rural areas as a covariate. The sociodemographic factors were selected based on the previous studies that focused on the BMI of Bangladeshi women, particularly using the BDHS<sup>(52–55)</sup>.

Age was a continuous variable, whereas parity (0, 1, 2, 3,  $\geq 4$ ), education (none, primary, secondary, higher), wealth index (poorest, poorer, middle, richer, richest), division (Barisal, Chittagong, Dhaka, Khulna, Rajshahi, Sylhet), marital status (married, not married (i.e. divorced, widowed, separated, not living together)) and residence (urban, rural) were categorical variables. Rajshahi division was separated into two divisions (Rangpur and Rajshahi) in 2010. Hence, seven divisions were listed in BDHS 2011 and 2014. However, we merged these two into one division (Rajshahi) in the combined data set to be consistent with BDHS 2004 and 2007. The DHS created a household wealth index with five groups corresponding to poorest, poorer, middle, richer and richest from the reported households’ asset variables using principal component analysis<sup>(56)</sup>.


### Ethical approval

The present paper does not contain any studies with human participants performed by any of the authors. The

**Table 1** Frequency distribution of household sociodemographic characteristics and BMI of women of reproductive age (15–49 years, *n* 53 077), Bangladesh Demographic and Health Surveys (BDHS), 2004–2014

Sociodemographic factor	BDHS 2004		BDHS 2007		BDHS 2011		BDHS 2014	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
Age† (years), mean	30.45	–	31.08	–	31.25	–	31.45	–
Parity								
0	1034	9.9	902	9.1	1439	8.9	1462	8.8
1	1761	16.9	1865	18.8	3136	19.4	3481	21.0
2	2151	20.6	2258	22.7	4124	25.5	4451	26.9
3	1798	17.2	1713	17.2	3159	19.6	3213	19.4
$\geq 4$	3688	35.4	3208	32.3	4288	26.6	3646	23.8
Education								
No education	4180	40.1	3305	33.2	4379	27.1	4059	24.5
Primary	3105	29.8	3002	30.2	4911	30.4	4909	29.7
Secondary	2564	24.6	2880	29.0	5633	34.9	6083	36.7
Higher	583	5.6	759	7.6	1223	7.6	1502	9.1
Wealth index								
Poorest	1924	18.4	1657	16.7	2875	17.8	3055	18.5
Poorer	1921	18.4	1810	18.2	3025	18.7	3103	18.7
Middle	1952	18.7	1887	19.0	3088	19.1	3340	20.2
Richer	2027	19.4	1973	19.8	3395	21.0	3496	21.1
Richest	2608	25.0	2619	26.3	3763	23.3	3559	21.5
Division								
Barisal	1220	11.7	1297	13.0	1797	11.1	1961	11.8
Chittagong	1836	17.6	1732	17.4	2575	15.9	2617	15.8
Dhaka	2380	22.8	2119	21.3	2791	17.3	2875	17.4
Khulna	1601	15.3	1566	15.7	2418	15.0	2423	14.6
Rajshahi	2359	22.6	1890	19.0	4668	28.9	4691	28.3
Sylhet	1036	9.9	1342	13.5	1897	11.7	1986	12.0
Marital status								
Married	9603	92.1	9170	92.2	15 099	93.5	15 580	94.1
Not married	829	7.9	776	7.8	1047	6.5	973	5.9
Residence								
Rural	6864	65.8	6168	62.0	10 510	65.1	10 824	65.4
Urban	3568	34.2	3778	38.0	5636	34.9	5729	34.6
BMI category								
Underweight	3405	32.6	2758	27.7	3967	22.9	3006	18.2
Normal	5910	56.7	5766	58.0	9490	58.8	9393	56.7
Overweight and obese	1117	10.7	1422	14.3	2959	18.9	414	25.1

†Continuous variable.



**Fig. 1** (colour online) The proportion of underweight (■), normal (■) and overweight (■), by urban–rural residence and year, among women of reproductive age (15–49 years,  $n$  53 077), Bangladesh Demographic and Health Surveys, 2004–2014

Bangladesh Medical Research Council and the Institutional Review Boards of ICF International approved the BDHS protocols, questionnaires and verbal consent forms for the respective years. All respondents were de-identified before publishing the data. The secondary data sets analysed during the current study are freely available upon request from the DHS website (<https://dhsprogram.com/data/available-datasets.cfm>).

## Results

Over the 10-year study period, the proportion of overweight women increased and the proportion of underweight women decreased (Fig. 1). The percentage of overweight increased during 2004–2014 from 10.7 to 25.1% (Table 1). In that period, the underweight percentage decreased almost by half, from 32.6 to 18.2%.

The average age of women in the last four surveys were similar, about 31 years (Table 1). Households with one or two children increased, whereas those with four or more children decreased. The proportion of ‘not married’ women in the four samples was 6–8% and the major portion of each sample (approximately 60%) was from rural areas. The percentage of uneducated women decreased from 40.1% (2004) to 24.5% (2014). As the wealth index was a simulated variable that classified household possessions into five levels, its values were proportionally equal for each survey.

The proportion of underweight women was much lower and the proportion of overweight women was nearly six times higher for the richest households (38.1%) than the poorest households (5.9%; Table 2). Only about 9%

of highly educated women were underweight compared with 32.3% of uneducated women. In contrast, only about 11% of the women with no education were overweight, whereas 38.8% of the highly educated women were overweight in the combined sample. The highest percentages of overweight women were in Dhaka and Khulna divisions and the highest proportion of underweight women was from Sylhet division. Only 12.5% of residents in the rural areas were overweight compared with 28.8% in the urban areas. All of the categorical variables showed a significant association with underweight and overweight categories in the bivariate analysis  $\chi^2$  test (Table 2).

The generalized linear mixed-effect model fitted to the underweight category compared with the other BMI categories in both urban and rural areas showed a significant ( $P < 0.01$ ) trend over the years. In comparison to 2004, women living in urban areas of Bangladesh had 54% less chance of being underweight in 2014, when adjusted for other covariates (Table 3). In the combined model, where both urban and rural residents were studied, we reached a similar conclusion with women becoming less prone to underweight as the years have gone by. Age, education, wealth index, division and marital status showed a significant association with women’s underweight status. Highly educated women were half as likely to be underweight compared with the illiterate, and women belonging to the richest class had 75% less chance of being underweight compared with those in the poorest households. Compared with Barisal division, women in Sylhet were 1.4 times more likely to be underweight in both urban and rural areas. As per Table 3, there was hardly any difference between urban and rural areas in the magnitude of trend in their underweight status. However, the combined model found that

**Table 2** Bivariate analysis of BMI category by household sociodemographic characteristics of women of reproductive age (15–49 years, *n* 53 077) in four combined Bangladesh Demographic and Health Surveys, 2004–2014

Sociodemographic factor	BMI category						<i>P</i> value†
	Underweight ( $<18.50$ kg/m <sup>2</sup> )		Normal ( $18.50$ – $24.99$ kg/m <sup>2</sup> )		Overweight and obese ( $\geq 25.00$ kg/m <sup>2</sup> )		
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	
Age‡ (years), mean	30.45	–	31.08	–	31.25	–	<0.001
Parity							
0	1222	25.2	3015	62.3	600	12.4	<0.001
1	2567	25.1	5922	57.8	1754	17.2	
2	2501	19.3	7548	58.1	2935	22.6	
3	2122	21.5	5695	57.6	2065	20.1	
≥4	4454	29.4	8378	55.4	2298	15.2	
Year							
2004	3405	32.6	5910	56.7	1117	10.7	<0.001
2007	2758	27.7	5766	58.0	1422	14.3	
2011	3697	22.9	9490	58.8	2959	18.1	
2014	3006	18.2	9393	56.7	4144	25.1	
Education							
No education	5145	32.3	9081	57.0	1697	10.6	<0.001
Primary	4124	25.9	9378	58.9	2425	15.2	
Secondary	3240	18.9	9967	58.1	3953	23.0	
Higher	357	8.8	2133	52.4	1577	38.8	
Wealth index							
Poorest	3677	38.7	5268	55.4	563	5.9	<0.001
Poorer	3044	30.9	6022	61.1	796	8.1	
Middle	2664	25.9	6317	61.5	1286	12.1	
Richer	2199	20.2	6549	60.1	2143	19.7	
Richest	1282	10.2	6403	51.0	4864	38.1	
Division							
Barisal	1658	26.4	3649	58.2	968	15.1	<0.001
Chittagong	1934	22.1	5093	58.1	1733	19.8	
Dhaka	2354	23.2	5741	56.5	2070	20.4	
Khulna	1566	19.6	4745	59.3	1697	21.2	
Rajshahi	3267	24.0	8062	59.2	2279	16.8	
Sylhet	2087	33.3	3269	52.2	905	14.2	
Marital status							
Married	11 644	23.5	28 671	58.0	9137	18.4	<0.001
Not married	1222	33.7	1888	52.1	515	14.1	
Residence							
Rural	9689	28.2	20 414	59.4	4263	12.5	<0.001
Urban	3177	17.0	10 145	54.2	5389	28.8	

†*P* value of the  $\chi^2$  test between BMI and sociodemographic factors after adjusting the complex survey design (survey weights and cluster-wise variation) using second-order Rao–Scott corrections.

‡Continuous variable.

the urban women were 17% less likely to be underweight than the rural residents.

The trend in the overweight category was different for urban and rural areas (Table 4). In the urban model, the urban residents were over 2.4 times more likely to be overweight in 2014 compared with 2004. However, the magnitude of change in odds was higher in rural areas, as the women living there were over 3.9 times more likely to be overweight in 2014 compared with 2004 (rural model, Table 4). This indicates that the extent of nutritional shift was greater for rural residents. Age, parity, education, wealth index and marital status played a significant role in these models. The highly educated women were 2.35 (urban) and 1.75 times (rural) more likely to be overweight in 2014 compared with 2004. Similarly, the wealthier the household, the more likely (OR=7.25 for urban and

OR=5.66 for rural) that a woman from that household would be overweight compared with those from the poorer quintiles. The non-married (e.g. divorced, widowed or separated) women were less likely to be overweight in reference to the married women. Although the volume of shift was higher for the rural women, the urban residents still showed 50% more chance of being overweight, compared with the rural women, in 2014 *v.* 2004 (combined model, Table 4).

## Discussion

The current study demonstrates that there has been a substantial change in the BMI status of reproductive-age women in Bangladesh. Between 2004 and 2014, there

**Table 3** Binary generalized linear mixed-effect models for the association of underweight (*v.* normal/overweight/obese; reference) with household sociodemographic characteristics, according to urban–rural residence and overall, among women of reproductive age (15–49 years, *n* 53 077), Bangladesh Demographic and Health Surveys, 2004–2014

Sociodemographic factor	Urban residents		Rural residents		Combined	
	OR	95 % CI	OR	95 % CI	OR	95 % CI
Random effect (cluster) variance	1.09	–	1.09	–	1.09	–
Age (scaled)†	0.24	0.19, 0.30*	0.64	0.55, 0.73*	0.48	0.42, 0.54*
Parity						
0	1.00	Ref.	1.00	Ref.	1.00	Ref.
1	0.95	1.05, 0.87	1.20	1.09, 1.33*	1.11	1.02, 1.21*
2	0.70	0.60, 0.82*	0.89	0.80, 0.99*	0.81	0.75, 0.89*
3	0.79	0.66, 0.94*	0.94	0.84, 1.05	0.88	0.80, 0.97*
≥4	1.25	1.04, 1.51*	1.20	1.06, 1.34*	1.19	1.08, 1.32*
Year						
2004	1.00	Ref.	1.00	Ref.	1.00	Ref.
2007	0.83	0.72, 0.95*	0.81	0.74, 0.89*	0.81	0.75, 0.88*
2011	0.59	0.52, 0.68*	0.64	0.59, 0.70*	0.63	0.59, 0.68*
2014	0.46	0.40, 0.53*	0.46	0.42, 0.50*	0.46	0.43, 0.50*
Education						
No education	1.00	Ref.	1.00	Ref.	1.00	Ref.
Primary	0.80	0.72, 0.90*	0.86	0.81, 0.91*	0.84	0.80, 0.89*
Secondary	0.61	0.54, 0.69*	0.80	0.74, 0.87*	0.73	0.69, 0.78*
Higher	0.37	0.30, 0.45*	0.63	0.53, 0.75*	0.49	0.43, 0.56*
Wealth index						
Poorest	1.00	Ref.	1.00	Ref.	1.00	Ref.
Poorer	0.75	0.63, 0.88*	0.74	0.69, 0.79*	0.75	0.70, 0.80*
Middle	0.55	0.47, 0.65*	0.63	0.58, 0.67*	0.62	0.58, 0.66*
Richer	0.50	0.43, 0.58*	0.46	0.42, 0.50*	0.49	0.45, 0.52*
Richest	0.24	0.21, 0.29*	0.26	0.23, 0.30*	0.25	0.23, 0.27*
Division						
Barisal	1.00	Ref.	1.00	Ref.	1.00	Ref.
Chittagong	0.95	0.79, 1.15	0.81	0.72, 0.90*	0.85	0.77, 0.93*
Dhaka	0.97	0.81, 1.16	0.93	0.83, 1.03*	0.93	0.85, 1.02
Khulna	0.99	0.82, 1.20	0.68	0.60, 0.76*	0.76	0.69, 0.84*
Rajshahi	1.02	0.86, 1.22	0.84	0.76, 0.92*	0.88	0.81, 0.96*
Sylhet	1.42	1.16, 1.74*	1.45	1.29, 1.63*	1.40	1.30, 1.59*
Marital status						
Married	1.00	Ref.	1.00	Ref.	1.00	Ref.
Not married	1.56	1.35, 1.80*	1.60	1.45, 1.77*	1.60	1.47, 1.73*
Residence						
Rural					1.00	Ref.
Urban					0.83	0.79, 0.89*

Ref., reference category.

\*OR and 95 % CI significant at 1 %.

†Age (15–49 years) was scaled (min = 0, mean = 0.5, max = 1) in the model convergence.

was a 15 % increase in the prevalence of overweight status and a similar decrease in underweight status. The reduction in underweight status was of similar magnitude in both urban and rural areas; whereas there was a greater relative change in overweight status in the rural areas, which is congruent with recent review findings<sup>(57)</sup>. The underweight prevalence in rural areas remained relatively high, as did the overweight prevalence in the urban residents. These findings of the shift of nutritional burden are an extension of previous findings depicting consistency with literature from Bangladesh<sup>(52,58,59)</sup>. Apart from the year-wise change and variation by urban–rural residence, the current study also finds that education, family income level, marital status of the women and geographic divisional locations played significant roles in the double burden emergence in Bangladesh.

Women's level of education was a significant predictor of nutritional shift. The higher rate of overweight in women with higher education is presumably because these women are more likely to be in sedentary jobs with less physically intensive activities such as construction works<sup>(52)</sup>. Occupational sitting time is high for educated persons, who prefer desk jobs, which in turn increases the risk of overweight and obesity<sup>(60)</sup>. On the other hand, the economic activities of illiterate women, who generally do not have adequate technical skills, are mostly confined to labour-intensive work<sup>(61,62)</sup>. However, the BDHS provides neither the daily physical activity counts of the respondents nor sitting times. Another possible explanation could be that the educated women have less time to prepare home-cooked meals and tend to use the more convenient options (processed foods) that could lead to increased

**Table 4** Binary generalized linear mixed-effect models for the association of overweight/obesity (v. normal/underweight; reference) with household sociodemographic characteristics, according to urban/rural residence and overall, among women of reproductive age (15–49 years, *n* 53 077), Bangladesh Demographic and Health Surveys, 2004–2014

Sociodemographic factors	Urban residents		Rural residents		Combined	
	OR	95 % CI	OR	95 % CI	OR	95 % CI
Random effect(cluster) variance	1.09	–	1.09	–	1.09	–
Age (scaled)†	8.08	6.65, 9.83*	5.30	4.37, 6.43*	6.70	5.85, 7.69*
Parity						
0	1.00	Ref.	1.00	Ref.	1.00	Ref.
1	1.26	1.08, 1.46*	1.17	0.99, 1.37	1.22	1.09, 1.36*
2	1.51	1.30, 1.75*	1.37	1.17, 1.61*	1.46	1.31, 1.63*
3	1.41	1.20, 1.67*	1.21	1.02, 1.43	1.33	1.18, 1.50*
≥4	1.16	0.97, 1.39	0.92	0.77, 1.10	1.04	0.91, 1.18
Year						
2004	1.00	Ref.	1.00	Ref.	1.00	Ref.
2007	1.18	1.02, 1.37	1.45	1.24, 1.69*	1.31	1.18, 1.46*
2011	1.55	1.36, 1.78*	2.30	2.01, 2.64*	1.89	1.71, 2.08*
2014	2.47	2.16, 2.83*	3.91	3.42, 4.47*	3.11	2.83, 3.43*
Education						
No education	1.00	Ref.	1.00	Ref.	1.00	Ref.
Primary	1.46	1.30, 1.64*	1.34	1.22, 1.48*	1.40	1.30, 1.51*
Secondary	2.00	1.78, 2.25*	1.59	1.42, 1.77*	1.82	1.68, 1.97*
Higher	2.35	2.05, 2.71*	1.75	1.47, 2.07*	2.13	1.92, 2.37*
Wealth index						
Poorest	1.00	Ref.	1.00	Ref.	1.00	Ref.
Poorer	1.41	1.07, 1.85	1.26	1.10, 1.43*	1.27	1.13, 1.42*
Middle	2.18	1.71, 2.78*	1.87	1.65, 2.12*	1.88	1.68, 2.10*
Richer	2.98	2.38, 3.75*	3.14	2.76, 3.56*	2.86	2.57, 3.19*
Richest	7.25	5.78, 9.09*	5.66	4.90, 6.53*	6.14	5.48, 6.88*
Division						
Barisal	1.00	Ref.	1.00	Ref.	1.00	Ref.
Chittagong	1.02	0.86, 1.22	1.42	1.22, 1.66*	1.24	1.10, 1.39*
Dhaka	0.93	0.79, 1.09	1.26	1.08, 1.48*	1.11	0.99, 1.25
Khulna	1.06	0.89, 1.26	1.52	1.30, 1.77*	1.31	1.16, 1.47*
Rajshahi	1.03	0.87, 1.21	1.16	1.00, 1.34	1.11	0.99, 1.24
Sylhet	0.81	0.67, 0.99	0.88	0.74, 1.05	0.86	0.75, 0.98
Marital status						
Married	1.00	Ref.	1.00	Ref.	1.00	Ref.
Not married	0.64	0.55, 0.74*	0.75	0.64, 0.88*	0.68	0.61, 0.76*
Residence						
Rural					1.00	Ref.
Urban					1.52	1.43, 1.63*

Ref., reference category.

\*OR and 95 % CI significant at 1 %.

†Age (15–49 years) was scaled (min = 0, mean = 0.5, max = 1) in the model convergence.

weight<sup>(63,64)</sup>. In the study period (2004–2014), the findings suggest that the secondary or higher educated women were over 1.5 times more likely to be overweight, whereas only 10.5 % of the illiterate women were overweight (Table 4).

Economic solvency affected the shift of BMI scales from underweight to overweight in Bangladesh as purchasing capacity of the general mass increased. The poor uneducated were mostly undernourished, while the rich educated were overweight. This leads to the assumption that women living in well-off families had access to surplus income and consequently increased their intake of nutritious (and/or processed) foods, whereas poor families could not afford them and remained underweight and undernourished<sup>(65)</sup>. Another possibility in the urban setting could be the overconsumption of fast (or junk) foods, which are more accessible to the wealthier sections of

society, that brings a change in the traditional dietary intake that elevates weight in general<sup>(66,67)</sup>. Thus, from the surveys between 2004 and 2014, it can be observed that 73 % of the overweight women were from above middle wealth quintiles.

From the divisional variation, it is eminent that underweight status increased significantly only in Sylhet compared with Barisal, and in other divisions the overall BMI increased with a significant decrease in underweight status. These divisions also experienced an increase in overweight status compared with Barisal division. This could be explained by the educational status and income of households, as Sylhet division lags behind in education and Barisal in economy<sup>(68–70)</sup>. Marital status in all models showed that women, who were not living with their partners and were divorced, seemed to be leaning towards underweight compared with currently married women.



This could be because childbirth leads to gain of weight (parity in Tables 3 and 4) and married women are less likely to care about their weight increase<sup>(71–73)</sup>. One other possibility could be that single women are relatively more economically disadvantaged than married ones where the husband contributes to income.

Urbanization creates a fast-paced economic paradigm where population density increases and food consumption follows the global trend, which currently refers to processed foods or ‘western’ diets more often filled with fats and refined carbohydrates<sup>(63,74,75)</sup>. In the current context of Bangladesh, various restaurants, food parks and supermarkets are gaining popularity in the urban areas, serving as places for recreational family activities, further fuelled by media publicity<sup>(76,77)</sup>. This changes the everyday food intake of the urban residents. Moreover, with continuous urban migration towards already crammed cities, increased housing and infrastructures have dried up the open spaces, which eventually results in an opportunity cost for the residents’ physical activities<sup>(78–80)</sup>. Thus, outdoor activity-based recreations are replaced by home-based entertainments (e.g. televisions, computers, board games), which ultimately lead to increased sitting time<sup>(81)</sup>. These could explain why the current study finds that, compared with rural residents, more than twice the proportion of overweight women were urban residents in Bangladesh.

Bangladesh has recently been classified as a low middle-income country following strong growth rate in gross domestic product in recent times, and it aims to become a middle-income country by 2021<sup>(82,83)</sup>. In the process of economic development, Bangladesh has shown commendable success in the public health sector, including rapid reduction of undernutrition<sup>(84,85)</sup>. Bangladesh’s Integrated Nutrition Program 1995–2004 and the subsequent National Nutritional Program had debatable, limited impact on existing nutritional status<sup>(86–89)</sup>. However, foreign investment paired with the efforts of local non-governmental organizations has improved maternal health care, which has reduced the underweight and undernutrition scenario in distant corners of Bangladesh, decentralizing the health benefits<sup>(85,90,91)</sup>. The findings of the present study support that the proportion of underweight women of reproductive age has declined over the last decade, even in the rural areas, to which growth in household economic status and increased female education have contributed.

This observed growth in gross domestic product was found to be associated with the double burden of malnutrition in recent literature, both in Asia and worldwide<sup>(92,93)</sup>. Economic growth has reached the rural areas with much developed road infrastructure and that brought application of commercial products for agriculture and day-to-day household tasks<sup>(57)</sup>. This has led to a decreased need for manual labour and with the additional availability of processed carbohydrates all over the country, to higher energy intake<sup>(94,95)</sup>. Although our discussion is focused on urban and rural differences, previous studies on nutrition in

Bangladesh have found differences between adult men and women<sup>(59)</sup>. It has already been observed that such double burden is affecting children (e.g. stunting) in low- and middle-income countries<sup>(96)</sup>. Thus, economic development demands a revisit to the existing nutritional policies to address the double burden of malnutrition.

While the current health policies focusing on nutrition generally emphasize the cases of undernutrition and underweight, multifaceted national policies from evidence-based experimental interventions are required to address the double nutritional burden<sup>(85,97)</sup>. Similarly, the UN Decade of Action on Nutrition has already addressed the double burden as an emerging problem worldwide and called for cross-cutting and coherent policies and programmes<sup>(98)</sup>. Such intervention strategies are tricky given that both ends of the nutritional spectrum are affected by the same covariates with a varied range. For example, the wealth index was found to be a significant factor for both nutrition statuses; however, the richest households are more likely to be overweight and poor households tend to be underweight. As a part of the UN Decade of Action on Nutrition, in 2016 the WHO suggested six areas of policy action including promoting sustainable diets by encouraging local production, setting up social protection programmes with awareness campaigns, and applying national cross-government, multisectoral, multi-stakeholder mechanisms to incorporate international guidelines on healthy diets in local communities<sup>(99)</sup>. However, all these would require intervention pilot studies in the context of Bangladesh prior to national policy implementation.

There were some limitations to the present study. First, there is no particular set of BMI levels for women in Bangladesh, which undermines the proper weight/height quantification as BMI tends to vary with a number of factors: racial group, ethnicity, geographic location, gender, age, hormone levels and other socio-economic measures<sup>(100–102)</sup>. More importantly, the typical threshold for overweight ( $\geq 25.00$  kg/m<sup>2</sup>) might underestimate the current problem of overnutrition in Bangladesh<sup>(103)</sup>. This could be one of the reasons why we found only 0.1% of obese women in the study period. Second, the study was limited to females in their reproductive age. Hence, generalization of the results and interpretation of the findings must be cautiously done. Third, respondents’ food intake, physical condition and sitting time, as well as national-level factors such as food prices and quality of nutrition are important components while analysing nutritional status, which were not available in BDHS data sets. Fourth, the sample size varied across the data sets over the years, which might compromise the interpretation across surveys; however, consistent methodologies were followed in all the surveys. Finally, the trend analysis would be more substantial with longitudinal data, particularly if the same respondents could be followed from rural–urban migration to assess their changes. Future studies could consider these issues and adjust the models for further precise findings.







## Conclusions

Bangladesh is transitioning from a low-income to a middle-income country, and its sociodemographic changes are influencing the public health sector. Generally, in Bangladesh, the primary concern for maternal health has been undernourishment and underweight. However, the current study argues that the nutrition scale is shifting; overall, the prevalence of underweight status is decreasing and there has been a consequent increase of overweight status. Prevalence of underweight status has remained high in the rural areas and the prevalence of overweight is increasing rapidly in both rural and urban areas. The significant contributors to this double burden were the change in women's level of education, increased household wealth, divisional location and rapid urbanization. Women with higher education from well-off families living in urban areas seem to be overweight, whereas uneducated women from poor households residing in the rural areas are underweight. To characterize these changes in the context of Bangladesh, health policies focusing only on undernutrition need to be revised and consider the new double burden. Revised policies, however, would require application of multiple area-based intervention studies in line with the UN Decade of Action on Nutrition to address the double nutritional burden in Bangladesh.

## Acknowledgements

**Financial support:** This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors. **Conflict of interest:** There is no conflict of interest among the authors. **Authorship:** R.K.B. conceptualized the study, compiled the data, synthesized the analysis plan and drafted the manuscript. N.R. conducted the statistical analysis and interpreted the findings. R.K. and A.H.B. provided expert opinions and critically reviewed the manuscript. S.A. overviewed the statistical analysis and revised the manuscript. All authors read and approved the manuscript. **Ethics of human subject participation:** This article does not contain any studies with human participants performed by any of the authors. The Bangladesh Medical Research Council and the Institutional Review Boards of ICF International approved the BDHS protocols, questionnaires and verbal consent forms for the respective years. All respondents were de-identified before publishing the data. The secondary data sets analysed during the current study are freely available upon request from the DHS website (<https://dhsprogram.com/data/available-datasets.cfm>).

## References

1. Finucane MM, Stevens GA, Cowan MJ *et al.* (2011) National, regional, and global trends in body-mass index since 1980: systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9.1 million participants. *Lancet* **377**, 557–567.
2. Prentice AM (2005) The emerging epidemic of obesity in developing countries. *Int J Epidemiol* **35**, 93–99.
3. Popkin BM, Adair LS & Ng SW (2012) Global nutrition transition and the pandemic of obesity in developing countries. *Nutr Rev* **70**, 3–21.
4. Loret de Mola C, Quispe R, Valle GA *et al.* (2014) Nutritional transition in children under five years and women of reproductive age: a 15-years trend analysis in Peru. *PLoS One* **9**, e92550.
5. Abdullah A (2015) The double burden of undernutrition and overnutrition in developing countries: an update. *Curr Obes Rep* **4**, 337–349.
6. Shrimpton R, Mbuya NV & Provo AM (2016) *The Double Burden of Malnutrition in East Asia and the Pacific: Evidence and Lessons for a Multisectoral Response. Health, Nutrition, and Population (HNP) Discussion Paper*. Washington, DC: World Bank Group; available at <http://documents.worldbank.org/curated/en/185291487250172586/The-double-burden-of-malnutrition-in-East-Asia-and-the-Pacific-evidence-and-lessons-for-a-multisectoral-response> (accessed September 2019).
7. Barquera S, Pedroza-Tobias A & Medina C (2016) Cardiovascular diseases in mega-countries: the challenges of the nutrition, physical activity and epidemiologic transitions, and the double burden of disease. *Curr Opin Lipidol* **27**, 329–344.
8. Khan SH & Talukder SH (2013) Nutrition transition in Bangladesh: is the country ready for this double burden. *Obes Rev* **14**, Suppl. 2, 126–133.
9. Khanam R, Lee ASC, Ram M *et al.* (2018) Levels and correlates of nutritional status of women of childbearing age in rural Bangladesh. *Public Health Nutr* **21**, 3037–3047.
10. Nawaz S (2010) Microfinance and poverty reduction: evidence from a village study in Bangladesh. *J Asian Afr Stud* **45**, 670–683.
11. Davis P & Baulch B (2011) Parallel realities: exploring poverty dynamics using mixed methods in rural Bangladesh. *J Dev Stud* **47**, 118–142.
12. Ahmed T, Roy S, Alam N *et al.* (2012) Determinants of undernutrition in children under 2 years of age from rural Bangladesh. *Indian Pediatr* **49**, 821–824.
13. Chowdhury AM, Bhuiya A, Chowdhury ME *et al.* (2013) The Bangladesh paradox: exceptional health achievement despite economic poverty. *Lancet* **382**, 1734–1745.
14. Das SK, Chisti MJ, Malek MA *et al.* (2015) Changing childhood malnutrition in Bangladesh: trends over the last two decades in urban–rural differentials (1993–2012). *Public Health Nutr* **18**, 1718–1727.
15. National Institute of Population Research and Training, Mitra and Associates & ICF International (2016) *Bangladesh Demographic and Health Survey 2014*. Dhaka and Rockville, MD: NIPORT, Mitra and Associates, and ICF International.
16. Vos T, Allen C, Arora M *et al.* (2016) Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* **388**, 1545–1602.
17. Arifeen SE, Hill K, Ahsan KZ *et al.* (2014) Maternal mortality in Bangladesh: a Countdown to 2015 country case study. *Lancet* **384**, 1366–1374.
18. Liu L, Oza S, Hogan D *et al.* (2016) Global, regional, and national causes of under-5 mortality in 2000–15: an updated systematic analysis with implications for the Sustainable Development Goals. *Lancet* **388**, 3027–3035.
19. Akhter S & Dasvarma G (2017) Whither MDG 5 in Bangladesh and its regions? *J Popul Res* **34**, 279–301.



20. Islam MM, Khan AM & Islam MM (2013) Textile industries in Bangladesh and challenges of growth. *Res J Eng Sci* **2278**, 9472.
21. Gautam M & Faruque R (2016) *Dynamics of Rural Growth in Bangladesh: Sustaining Poverty Reduction*. Washington, DC: World Bank Publications.
22. White H (2005) *Maintaining Momentum to 2015? An Impact Evaluation of Interventions to Improve Maternal and Child Health and Nutrition in Bangladesh*. Washington, DC: World Bank Publications.
23. Nahar S, Mascie-Taylor CG & Begum HA (2009) Impact of targeted food supplementation on pregnancy weight gain and birth weight in rural Bangladesh: an assessment of the Bangladesh Integrated Nutrition Program (BINP). *Public Health Nutr* **12**, 1205–1212.
24. Belton B, Mariana van Asseldonk IJ & Thilsted SH (2014) Faltering fisheries and ascendant aquaculture: implications for food and nutrition security in Bangladesh. *Food Policy* **44**, 77–87.
25. Subramanian SV, Perkins JM, Ozaltin E *et al.* (2011) Weight of nations: a socioeconomic analysis of women in low- to middle-income countries. *Am J Clin Nutr* **93**, 413–421.
26. Jones-Smith JC, Gordon-Larsen P, Siddiqi A *et al.* (2012) Is the burden of overweight shifting to the poor across the globe? Time trends among women in 39 low- and middle-income countries (1991–2008). *Int J Obes (Lond)* **36**, 1114–1120.
27. Black RE, Victora CG, Walker SP *et al.* (2013) Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet* **382**, 427–451.
28. Barrett CB (2010) Measuring food insecurity. *Science* **327**, 825–828.
29. Ha DTP, Feskens EJM, Deurenberg P *et al.* (2011) Nationwide shifts in the double burden of overweight and underweight in Vietnamese adults in 2000 and 2005: two national nutrition surveys. *BMC Public Health* **11**, 62.
30. Ruel MT & Alderman H; Maternal and Child Nutrition Study Group (2013) Nutrition-sensitive interventions and programmes: how can they help to accelerate progress in improving maternal and child nutrition? *Lancet* **382**, 536–551.
31. Headey DD (2013) Developmental drivers of nutritional change: a cross-country analysis. *World Dev* **42**, 76–88.
32. Liu S, Allen A & Fraser W (2008) Fetal and infant health outcomes. In *Canadian Perinatal Health Report 2008 Edition*, p. 123. Ottawa: Public Health Agency of Canada.
33. Han Z, Mulla S, Beyene J *et al.* (2010) Maternal underweight and the risk of preterm birth and low birth weight: a systematic review and meta-analyses. *Int J Epidemiol* **40**, 65–101.
34. Ozaltin E, Hill K & Subramanian SV (2010) Association of maternal stature with offspring mortality, underweight, and stunting in low- to middle-income countries. *JAMA* **303**, 1507–1516.
35. Khanam R, Lee ACC, Mitra DK *et al.* (2019) Maternal short stature and underweight status are independent risk factors for preterm birth and small for gestational age in rural Bangladesh. *Eur J Clin Nutr* **73**, 733–742.
36. Ahmed T, Hossain M & Sanin KI (2012) Global burden of maternal and child undernutrition and micronutrient deficiencies. *Ann Nutr Metab* **61**, Suppl. 1, 8–17.
37. Jaacks LM, Slining MM & Popkin BM (2015) Recent underweight and overweight trends by rural–urban residence among women in low- and middle-income countries. *J Nutr* **145**, 352–357.
38. Ahmed T, Mahfuz M, Ireen S *et al.* (2012) Nutrition of children and women in Bangladesh: trends and directions for the future. *J Health Popul Nutr* **30**, 1–11.
39. Ng SW, Zaghoul S, Ali HI *et al.* (2011) The prevalence and trends of overweight, obesity and nutrition-related non-communicable diseases in the Arabian gulf states. *Obes Rev* **12**, 1–13.
40. Ng M, Fleming T, Robinson M *et al.* (2014) Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* **384**, 766–781.
41. Look AHEAD Research Group, Gregg EW, Jakicic JM *et al.* (2016) Association of the magnitude of weight loss and changes in physical fitness with long-term cardiovascular disease outcomes in overweight or obese people with type 2 diabetes: a post-hoc analysis of the Look AHEAD randomised clinical trial. *Lancet Diabetes Endocrinol* **4**, 913–921.
42. Lavie CJ, Schutter AD, Parto P *et al.* (2016) Obesity and prevalence of cardiovascular diseases and prognosis – the obesity paradox updated. *Prog Cardiovasc Dis* **58**, 537–547.
43. Basu S, McKee M, Galea G *et al.* (2013) Relationship of soft drink consumption to global overweight, obesity, and diabetes: a cross-national analysis of 75 countries. *Am J Public Health* **103**, 2071–2077.
44. Neuman M, Kawachi I, Gortmaker S *et al.* (2013) Urban–rural differences in BMI in low- and middle-income countries: the role of socioeconomic status. *Am J Clin Nutr* **97**, 428–436.
45. Biswas T, Uddin MJ, Mamun AA *et al.* (2017) Increasing prevalence of overweight and obesity in Bangladeshi women of reproductive age: findings from 2004 to 2014. *PLoS One* **12**, e0181080.
46. Kamal SM, Hassan CH & Alam GM (2015) Dual burden of underweight and overweight among women in Bangladesh: patterns, prevalence, and sociodemographic correlates. *J Health Popul Nutr* **33**, 92–105.
47. Rutstein SO, Johnson K & ORC Macro MEASURE *et al.* (2004) *The DHS Wealth Index*. Calverton, MD: ORC Macro, MEASURE DHS.
48. World Health Organization (2000) *Obesity: Preventing and Managing the Global Epidemic. Report of a WHO Consultation. WHO Technical Report Series* no. 894. Geneva: WHO.
49. Bangladesh Bureau of Statistics (2015) *Population Monographs of Bangladesh: Age–Sex Composition of Bangladesh Population*. Dhaka: BBS, Statistics and Informatics Division, Ministry of Planning.
50. Thomas DR & Decady YJ (2004) Testing for association using multiple response survey data: approximate procedures based on the Rao–Scott approach. *Int J Test* **4**, 43–59.
51. Rabe-Hesketh S & Skrondal A (2008) Generalized linear mixed-effects models. In *Longitudinal Data Analysis: A Handbook of Modern Statistical Methods*, pp. 79–106 [G Fitzmaurice, M Davidian, G Verbeke *et al.*, editors]. Boca Raton, FL: Chapman & Hall/CRC.
52. Balarajan Y & Villamor E (2009) Nationally representative surveys show recent increases in the prevalence of overweight and obesity among women of reproductive age in Bangladesh, Nepal, and India. *J Nutr* **139**, 2139–2144.
53. Mostafa Kamal SM & Islam A (2010) Socio-economic correlates of malnutrition among married women in Bangladesh. *Malays J Nutr* **16**, 349–359.
54. Corsi DJ, Kyu HH & Subramanian SV (2011) Socioeconomic and geographic patterning of under- and overnutrition among women in Bangladesh. *J Nutr* **141**, 631–638.
55. Mohsena M, Goto R & Mascie-Taylor CN (2016) Maternal nutritional status (as measured by height, weight and BMI) in Bangladesh: trends and socio-economic association over the period 1996 to 2007. *Public Health Nutr* **19**, 1438–1445.
56. Howe LD, Hargreaves JR, Gabrysch S *et al.* (2009) Is the wealth index a proxy for consumption expenditure? A systematic review. *J Epidemiol Community Health* **63**, 871–877.



57. NCD Risk Factor Collaboration *et al.* (2019) Rising rural body-mass index is the main driver of the global obesity epidemic in adults. *Nature* **569**, 260–264.
58. Khan MM & Kraemer A (2009) Factors associated with being underweight, overweight and obese among ever-married non-pregnant urban women in Bangladesh. *Singapore Med J* **50**, 804–813.
59. Hoque ME, Hasan MT, Rahman M *et al.* (2017) Double burden of underweight and overweight among Bangladeshi adults differs between men and women: evidence from a nationally representative survey. *Public Health Nutr* **20**, 2183–2191.
60. Mummery WK, Schofield GM, Steele R *et al.* (2005) Occupational sitting time and overweight and obesity in Australian workers. *Am J Prev Med* **29**, 91–97.
61. Kabeer N, Mahmud S & Tasneem S (2011) *Does Paid Work Provide a Pathway to Women's Empowerment? Empirical Findings from Bangladesh*. Brighton: Institute of Development Studies.
62. Kumar D, Hossain A & Gope MC (2015) Role of micro credit program in empowering rural women in Bangladesh: a study on Grameen Bank Bangladesh Limited. *Asian Bus Rev* **3**, 114–120.
63. Schmidhuber J & Shetty P (2005) The nutrition transition to 2030. Why developing countries are likely to bear the major burden. *Acta Agric Scand Sec C* **2**, 150–166.
64. Baker P & Friel S (2014) Processed foods and the nutrition transition: evidence from Asia. *Obes Rev* **15**, 564–577.
65. Das S, Fahim SM, Islam MS *et al.* (2019) Prevalence and sociodemographic determinants of household-level double burden of malnutrition in Bangladesh. *Public Health Nutr* **22**, 1425–1432.
66. Rahman S (2014) Obesity in junk food generation in Asia: a health time bomb that needs early defusing! *South East Asia J Public Health* **3**, 1–2.
67. Morgan K & Sonnino R (2010) The urban foodscape: world cities and the new food equation. *Camb J Regions Econ Soc* **3**, 209–224.
68. Islam N, Angeles G, Mahub A *et al.* (2006) *Slums of Urban Bangladesh: Mapping and Census 2005*. Dhaka and Chapel Hill, NC: Centre for Urban Studies, National Institute of Population Research and Training, and MEASURE Evaluation.
69. Kabir I, Khanam M, Agho KE *et al.* (2012) Determinants of inappropriate complementary feeding practices in infant and young children in Bangladesh: secondary data analysis of demographic health survey 2007. *Matern Child Nutr* **8**, Suppl. 1, 11–27.
70. Tareque MI, Begum S & Saito Y (2014) Inequality in disability in Bangladesh. *PLoS One* **9**, e103681.
71. Kutob RM, Yuan NP, Wertheim BC *et al.* (2017) Relationship between marital transitions, health behaviors, and health indicators of postmenopausal women: results from the Women's Health Initiative. *J Womens Health* **26**, 313–320.
72. Biswas RK, Kabir E & Khan HT (2019) Socioeconomic transition and its influence on body mass index (BMI) pattern in Bangladesh. *J Eval Clin Pract* **25**, 130–141.
73. Chowdhury MAB, Adnan MM & Hassan MZ (2018) Trends, prevalence and risk factors of overweight and obesity among women of reproductive age in Bangladesh: a pooled analysis of five national cross-sectional surveys. *BMJ Open* **8**, e018468.
74. Popkin BM (2014) Nutrition, agriculture and the global food system in low and middle income countries. *Food Policy* **47**, 91–96.
75. Neiderud CJ (2015) How urbanization affects the epidemiology of emerging infectious diseases. *Infect Ecol Epidemiol* **5**, 27060.
76. Harun MA, Ahmed F & Maniruzzaman (2013) Customer hospitality: the case of fast food industry in Bangladesh. *World J Soc Sci* **3**, 88–104.
77. Ashraf MA, Akhter S & Noor SI (2014) Consumer behavior in fast food marketing in Bangladesh: a case study. *Develop Countr Stud* **4**, 34–44.
78. Reardon T, Boughton D, Tschirley D *et al.* (2015) Urbanization, diet change, and transformation of the downstream and midstream of the agrifood system: effects on the poor in Africa and Asia. *Faith Econ* **66**, 43–63.
79. Neupane S, Prakash KC & Doku DT (2016) Overweight and obesity among women: analysis of demographic and health survey data from 32 sub-Saharan African countries. *BMC Public Health* **16**, 30.
80. Melby CL, Orozco F, Ochoa D *et al.* (2017) Nutrition and physical activity transitions in the Ecuadorian Andes: differences among urban and rural dwelling women. *Am J Hum Biol* **29**, e22986.
81. Goryakin Y & Suhrcke M (2014) Economic development, urbanization, technological change and overweight: what do we learn from 244 Demographic and Health Surveys? *Econ Hum Biol* **14**, 109–127.
82. Feldman S (2015) Bangladesh in 2014. *Asian Surv* **55**, 67–74.
83. World Bank (2016) Helping Bangladesh Reach Middle Income Country Status. [http://www.worldbank.org/en/news/feature/2016/04/07/World\\_Bank\\_Group\\_s\\_New\\_Country\\_Partnership\\_Framework\\_helps\\_Bangladesh\\_Reach\\_Middle\\_Income\\_Country\\_Status](http://www.worldbank.org/en/news/feature/2016/04/07/World_Bank_Group_s_New_Country_Partnership_Framework_helps_Bangladesh_Reach_Middle_Income_Country_Status) (accessed November 2017).
84. Taylor L (2012) *The Nutrition Agenda in Bangladesh: 'Too Massive to Handle'? Analysing Nutrition Governance: Bangladesh Country Report*. Brighton: Institute of Development Studies.
85. Headey D, Hoddinott J, Ali D *et al.* (2015) The other Asian enigma: explaining the rapid reduction of undernutrition in Bangladesh. *World Dev* **66**, 749–761.
86. Hossain SM, Duffield A & Taylor A (2005) An evaluation of the impact of a US \$60 million nutrition programme in Bangladesh. *Health Policy Plan* **20**, 35–40.
87. Levinson FJ & Rohde JE (2005) Responses to: 'an evaluation of the impact of a US \$60 million nutrition programme in Bangladesh'. *Health Policy Plan* **20**, 405–406.
88. White H (2005) Comment on contributions regarding the impact of the Bangladesh integrated nutrition project. *Health Policy Plan* **20**, 408–411.
89. Sack DA, Roy SK, Ahmed T *et al.* (2005) An evaluation of the impact of a US \$60 million nutrition programme in Bangladesh (letter) (reply). *Health Policy Plan* **20**, 406–407.
90. Islam MS (2016) The NGOs sector in Bangladesh: emergence, contribution and current debate. *Adv Asian Soc Sci* **7**, 1182–1188.
91. Hasan MT, Soares Magalhaes RJ, Williams GM *et al.* (2016) Long-term changes in childhood malnutrition are associated with long-term changes in maternal BMI: evidence from Bangladesh, 1996–2011. *Am J Clin Nutr* **104**, 1121–1127.
92. Gillespie S & Haddad L (2001) *Attacking the Double Burden of Malnutrition in Asia and the Pacific*. Manila and Washington, DC: Asian Development Bank and International Food Policy Research Institute.
93. Kraak VI, Swinburn B, Lawrence M *et al.* (2011) The accountability of public–private partnerships with food, beverage and quick-serve restaurant companies to address global hunger and the double burden of malnutrition. *United Nations System Standing Committee on Nutrition: News* **39**, 11–24.
94. Reardon T, Timmer CP, Barrett CB *et al.* (2003) The rise of supermarkets in Africa, Asia, and Latin America. *Am J Agric Econ* **85**, 1140–1146.



95. Reardon T, Timmer CP & Minten B (2012) Supermarket revolution in Asia and emerging development strategies to include small farmers. *Proc Natl Acad Sci U S A* **109**, 12332–12337.
96. Tzioumis E, Kay MC, Bentley ME *et al.* (2016) Prevalence and trends in the childhood dual burden of malnutrition in low-and middle-income countries, 1990–2012. *Public Health Nutr* **19**, 1375–1388.
97. Motlagh ME, Kelishadi R, Amirkhani MA *et al.* (2011) Double burden of nutritional disorders in young Iranian children: findings of a nationwide screening survey. *Public Health Nutr* **14**, 605–610.
98. Demaio AR & Branca F (2018) Decade of action on nutrition: our window to act on the double burden of malnutrition. *BMJ Glob Health* **3**, Suppl. 1, e000492.
99. World Health Organization (2016) *The Double Burden of Malnutrition: Policy Brief*. Geneva: WHO.
100. Gray LJ, Yates T, Davies MJ *et al.* (2011) Defining obesity cut-off points for migrant South Asians. *PLoS One* **6**, e26464.
101. Bahk JY, Jung JH, Jin LM *et al.* (2010) Cut-off value of testes volume in young adults and correlation among testes volume, body mass index, hormonal level, and seminal profiles. *Urology* **75**, 1318–1323.
102. Bahat G, Tufan F, Saka B *et al.* (2012) Which body mass index (BMI) is better in the elderly for functional status? *Arch Gerontol Geriatr* **54**, 78–81.
103. Razak F, Anand SS, Shannon H *et al.* (2007) Defining obesity cut points in a multiethnic population. *Circulation* **115**, 2111–2118.